



TITLE:

On a family of subgroups of the Teichmuller modular group of genus two obtained from the Jones representation

AUTHOR(S):

MORISHITA, Masanori

CITATION:

MORISHITA, Masanori. On a family of subgroups of the Teichmuller modular group of genus two obtained from the Jones representation. 数理解析研究所講究録 1996, 962: 42-52

ISSUE DATE:

1996-08

URL:

<http://hdl.handle.net/2433/60546>

RIGHT:

On a family of subgroups of the Teichmüller modular group of genus two obtained from the Jones representation

Masanori MORISHITA

Department of Mathematics, Faculty of Science
Kanazawa University

Introduction

The purpose of the present paper is to give a family of “non-Torelli” subgroups of the Teichmüller modular group of genus 2 by confirming a conjecture, posed by Takayuki Oda, on the image of the Jones representation.

In [J], Jones attached to a Young diagram a Hecke algebra representation of the braid group B_n on n strings. As was shown in [ibid,10], the Jones representation of B_6 corresponding to the rectangular Young diagram $\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}$ factors through the Teichmüller modular group Γ of genus 2, namely, the mapping class group of a closed orientable surface of genus 2, and we thus get the representation $\pi : \Gamma \longrightarrow GL_5(\mathbf{Z}[x, x^{-1}])$ which is explicitly given ([ibid, p362]). Now, for a certain natural number n , specializing x to $\exp(2\pi\sqrt{-1}/n)$, we get a representation $\pi_n : \Gamma \longrightarrow GL_5(\mathcal{O}_K)$, where \mathcal{O}_K is the ring of integers in the n -th cyclotomic field K . Let F be the maximal real subfield of K and take a non-zero ideal I of \mathcal{O}_F , the ring of integers of F . The reduction of π_n modulo $I_K = I\mathcal{O}_K$ gives a representation $\pi_{n,I} : \Gamma \longrightarrow GL_5(\mathcal{O}_K/I_K)$. Then, Oda conjectured that the image of $\pi_{n,I}$ is a certain unitary group if I is prime to an ideal of \mathcal{O}_F containing (n) . (For the precise formulation, see Section 2).

The main result of this paper is to confirm Oda’s conjecture when I is a product of prime ideals of \mathcal{O}_F which are inert in K/F . The proof consists of two steps. We first show that $\pi_{n,\wp}$ is irreducible under certain conditions on n and a prime \wp , and then investigate the list of all irreducible subgroups of $PSL_5(\mathcal{O}_K/\wp_K)$ due to Martino and Wagoner [M-W]. For the case of a product of inert primes, we apply a criterion of Weisfeiler on the approximation of a Zariski-dense subgroup in a semisimple group over a

finite ring $[W]$. This proof is similar to that of Oda and Terasoma ([O-T]) for the similar problem on the Burau representations, where they use the induction after working with 2×2 matrices (see also [Be]). Our case is more complicated, for we work with 5×5 matrices and so the finite group theory is more involved.

We also check that the kernel of $\pi_{n,I}$ does not contain the Torelli group using its explicit generator given by Birman [B1].

Since the Teichmüller modular group is the fundamental group of the moduli space \mathcal{M} of compact Riemann surfaces of genus 2, our result gives a tower of 3-folds, namely, finite Galois coverings of \mathcal{M} with the Galois groups of finite unitary groups.

Notation. For an associative ring R with identity, $M_n(R)$ denotes the total matrix algebra over R of degree n , and $GL_n(R)$ denotes the groups of invertible elements of $M_n(R)$. We write R^\times for $GL_1(R)$. For $A \in M_n(R)$, tA , $tr(A)$, and $det(A)$ stand for the transpose, trace, and determinant of A , respectively. We write 0_n and 1_n for the zero and identity matrix in $M_n(R)$, respectively, and e_{ij} for the matrix unit and $diag(\cdot)$ for the diagonal matrix.

1. The Jones representation of the Teichmüller modular group of genus 2 and its unitarity

In [J], Jones attached to each Young diagram with n tiles a Hecke algebra representation of the braid group B_n on n strings. As was shown in [ibid, Section 10], the representation of B_6 corresponding to the rectangular Young diagram $\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}$ factors through the Teichmüller modular group Γ of genus 2, namely, the mapping class group of a closed orientable surface of genus 2. It is known that Γ admits the following presentation ([Bi2], Theorem 4.8, p 183-4).

generators: $\theta_1, \theta_2, \theta_3, \theta_4, \theta_5$.

defining relation:

$$\left\{ \begin{array}{l} \theta_i \theta_{i+1} \theta_i = \theta_{i+1} \theta_i \theta_{i+1} \quad (1 \leq i \leq 4), \\ \theta_i \theta_j = \theta_j \theta_i \quad (|i - j| \geq 2, 1 \leq i, j \leq 5), \\ (\theta_1 \theta_2 \theta_3 \theta_4 \theta_5)^6 = 1, \\ (\theta_1 \theta_2 \theta_3 \theta_4 \theta_5^2 \theta_4 \theta_3 \theta_2 \theta_1)^2 = 1, \\ \theta_1 \theta_2 \theta_3 \theta_4 \theta_5^2 \theta_4 \theta_3 \theta_2 \theta_1 \text{ commutes with } \theta_i \quad (1 \leq i \leq 5). \end{array} \right.$$

The Jones representation of Γ mentioned above is given explicitly on generators as follows ([J], p362).

$$\pi : \Gamma \longrightarrow GL_5(\mathbf{Z}[x, x^{-1}]), \quad x = t^{1/5};$$

$$\pi(\theta_1) = x^{-2} \begin{pmatrix} -1 & 0 & 0 & 0 & t \\ 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & t & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & t \end{pmatrix}, \quad \pi(\theta_2) = x^{-2} \begin{pmatrix} t & 0 & 0 & 0 & 0 \\ 0 & t & 0 & 0 & 0 \\ 0 & t & -1 & 0 & 0 \\ 1 & 0 & 0 & -1 & 0 \\ 1 & 0 & 0 & 0 & -1 \end{pmatrix},$$

$$\pi(\theta_3) = x^{-2} \begin{pmatrix} -1 & 0 & 0 & t & 0 \\ 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & t & 0 & 0 \\ 0 & 0 & 0 & t & 0 \\ 0 & 0 & 1 & 0 & -1 \end{pmatrix}, \quad \pi(\theta_4) = x^{-2} \begin{pmatrix} t & 0 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & t \\ 1 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & t \end{pmatrix},$$

$$\pi(\theta_5) = x^{-2} \begin{pmatrix} -1 & t & 0 & 0 & 0 \\ 0 & t & 0 & 0 & 0 \\ 0 & 0 & t & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & 0 & -1 \end{pmatrix}.$$

We see that $\det \pi(\theta_i) = -1, 1 \leq i \leq 5$.

Let $A = A(x) \in M_n(\mathbf{Z}[x, x^{-1}])$, $x = t^{1/5}$. We write A^* for ${}^tA(x^{-1})$ and call A x -hermitian if $A = A^*$. For a t -hermitian matrix A , we define the unitary group with respect to A by

$$U_n(A) := \{g \in GL_n(\mathbf{Z}[x, x^{-1}]) \mid g^* A g = A\}.$$

Lemma 1.1. *Let π be the representation given in Section 1. Then, there is a t -hermitian matrix $H \in M_5(\mathbf{Z}[x, x^{-1}])$ so that the image of π is contained in $U_5(H)$.*

Proof. By the straightforward computation, the following x -hermitian matrix satisfies the desired property.

$$\begin{pmatrix} (1+t)(1+t^{-1}) & -(1+t) & 2 & -(1+t) & -(1+t) \\ -(1+t^{-1}) & 1+t+t^{-1} & -(1+t^{-1}) & 1 & 1 \\ 2 & -(1+t) & (1+t)(1+t^{-1}) & -(1+t) & -(1+t) \\ -(1+t^{-1}) & 1 & -(1+t^{-1}) & 1+t+t^{-1} & 1 \\ -(1+t^{-1}) & 1 & -(1+t^{-1}) & 1 & 1+t+t^{-1} \end{pmatrix}.$$

If H' is such a matrix, then $H'H^{-1}$ commutes with $\pi(\theta_i)$, $1 \leq i \leq 5$. By the computation, we check that $H'H^{-1} \in \mathbf{Q}(x)^{\times} 1_5$.

We write $h = h_t$ for the matrix in the proof. We see that $\det(h_t) = (t + t^{-1})^4(1 + t + t^{-1})$.

2. The reduction of the specialized Jones representation at root of unity and the conjecture of Oda

Let n be a natural number. We assume that n is bigger than 2 and prime to 10. Let $\eta = \exp(2\pi\sqrt{-1}/n)$ and $\zeta = \eta^5$. Set $K = \mathbf{Q}(\zeta)$, $\mathcal{O}_K = \mathbf{Z}[\zeta]$, $F = \mathbf{Q}(\zeta + \zeta^{-1})$ and $\mathcal{O}_F = \mathbf{Z}[\zeta + \zeta^{-1}]$.

By specializing $t \rightarrow \zeta$, $x = t^{1/5} \rightarrow \eta$ in the representation π , we get a representation

$$\pi_n : \Gamma \longrightarrow GL_5(\mathcal{O}_K).$$

Take a non-zero ideal I of \mathcal{O}_F which is prime to n , and set $I_K = I\mathcal{O}_K$. The reduction of π_ζ modulo I_K defines the representation

$$\pi_{n,I} : \Gamma \longrightarrow GL_5(\mathcal{O}_K/I_K).$$

Then, $\pi_{n,I}$ certainly inherits the unitarity from π .

Lemma 2.1. *The image of $\pi_{n,I}$ is contained in*

$$U_5(\mathcal{O}_K/I_K; h_{n,I}) := \{g \in GL_5(\mathcal{O}_K/I_K) \mid g^* h_I g = h_I\},$$

where $h_{n,I} := h_\zeta \bmod I_K$ and $g^* = {}^t g^\tau$, τ is the involution induced from the generator of $\text{Gal}(K/F)$.

Proof. Immediate from Lemma 1.1. \square

To formulate the conjecture, we twist π_I a little bit. Let $\chi : \Gamma \rightarrow \mathcal{O}_K^\times$ be the character defined by $\chi(\theta_i) = -1$, and set $\chi_I := \chi \bmod I_K$. We then consider $\rho_I := \pi_{n,I} \otimes \chi_I$. Since $\det(\pi_\zeta(\theta_i)) = -1$, by Lemma 2.1, we have the inclusion

$$\rho_I(\Gamma) \subset SU_5(\mathcal{O}_K/I_K; h_{n,I}) := \{g \in U_5(\mathcal{O}_K/I_K; h_{n,I}) \mid \det(g) = 1\}.$$

Then, the conjecture posed by Oda is formulated as follows.

Conjecture 2.2. *There is a non-zero ideal \mathcal{C} of \mathcal{O}_F containing (n) so that the image of $\rho_{n,I}$ coincides with $SU_5(h_{n,I})$ if I is prime to \mathcal{C} .*

3. Non-split prime case

In this section, we verify Conjecture 2.2, when I is a maximal ideal \wp of \mathcal{O}_F which is inert in K/F . Set $\mathbf{F}_\wp = \mathcal{O}_F/\wp$, $\mathbf{F} = \mathbf{F}_{\wp K} = \mathcal{O}_K/\wp K$ for simplicity. We simply write π_\wp and ρ_\wp for $\pi_{n,\wp}$ and $\rho_{n,\wp}$, respectively, also h_\wp for $h_{n,\wp}$.

First, the following lemma shows each $\pi_\wp(\theta_i)$ is a quasi-reflection.

Lemma 3.1. *Assume that \wp is prime to $1 + \zeta$. Let $V = \mathbf{F}^{\oplus 5}$ be the representation space of π_\wp . For each $1 \leq i \leq 5$, there are subspaces X_i and Y_i of V such that*

$$\begin{aligned} V &= X_i \oplus Y_i, & \dim X_i &= 3, \dim Y_i = 2, \\ \pi_\wp(\theta_i)|_{X_i} &= -\eta^{-2} id_{X_i}, & \pi_\wp(\theta_i)|_{Y_i} &= \eta^3 id_{Y_i}, \end{aligned}$$

where η denotes a primitive n -th root of 1 in \mathbf{F} by abuse of notation.

Proof. By the direct computation, X_i and Y_i are given as follows:

$$\begin{aligned} X_1 &= \{ {}^t(x_1, x_2, 0, x_4, 0) \}, & Y_1 &= \{ {}^t(y_1, y_2, (1 + \zeta)y_2, y_2, (1 + \zeta^{-1})y_1) \} \\ X_2 &= \{ {}^t(0, 0, x_3, x_4, x_5) \}, & Y_2 &= \{ {}^t((1 + \zeta)y_1, (1 + \zeta^{-1})y_2, y_2, y_1, y_1) \} \\ X_3 &= \{ {}^t(x_1, x_2, 0, 0, x_5) \}, & Y_3 &= \{ {}^t(y_1, y_2, (1 + \zeta)y_2, (1 + \zeta^{-1})y_1, y_2) \} \\ X_4 &= \{ {}^t(0, x_2, x_3, x_4, 0) \}, & Y_4 &= \{ {}^t((1 + \zeta)y_1, y_1, y_2, y_1, (1 + \zeta^{-1})y_2) \} \\ X_5 &= \{ {}^t(x_1, 0, 0, x_4, x_5) \}, & Y_5 &= \{ {}^t(y_1, (1 + \zeta^{-1})y_1, (1 + \zeta)y_2, y_2, y_2) \}, \end{aligned}$$

where x_i 's and y_i 's run over \mathbf{F} and $\zeta = \eta^5$. \square

Lemma 3.2. Assume that \wp is prime to $(1 + \zeta)(\zeta + \zeta^{-1})(1 + \zeta + \zeta^{-1})$. Then, the representation π_\wp is irreducible.

Proof. Suppose that V has $\pi_\wp(\Gamma)$ -invariant subspace $W \neq 0, V$. First, assume $\dim(W) = 1$. Let w be a base of W and write $w = x + y$, $x \in X_1, y \in Y_1$. If $\pi_\wp(\theta_1)w = \alpha w$, $\alpha \in \mathbf{F}^\times$, by Lemma 4.1, we have $(\alpha + \eta^2)x + (\alpha - \eta^3)y = 0$, from which we see that $w \in X_1$ or $w \in Y_1$. Let $w = {}^t(x_1, x_2, 0, x_4, 0) \in X_1$. Then, $\pi_\wp(\theta_2)w = \eta^{-2} {}^t(\zeta x_1, \zeta x_2, \zeta x_2, x_1 - x_4, x_1)$ should be in X_1 and so we get $w = 0$. This is a contradiction. Similarly, w can not be in Y_1 . Hence, $\dim(W) > 1$. Note that the hermitian form $h_{n,\wp}$ is non-degenerate by our assumption. So, we may assume $\dim(W) = 2$, since the orthogonal complement of W with respect to $h_{n,\wp}$ is $\pi_\wp(\Gamma)$ -invariant. For this case, consider the exterior square representation $\wedge^2 \pi_\wp : \Gamma \longrightarrow GL(\wedge^2 V)$. Then, $\wedge^2 W$ is an invariant subspace of $\wedge^2 V$ and $\dim(\wedge^2 W) = 1$, and the similar argument to the above can be applied. Fix a basis of X_1 ; $v_1 = {}^t(1, 0, 0, 0, 0)$, $v_2 = {}^t(0, 1, 0, 0, 0)$, $v_3 = {}^t(0, 0, 0, 1, 0)$ and a basis of Y_1 ; $v_4 = {}^t(1, 0, 0, 0, 1 + \zeta^{-1})$, $v_5 = {}^t(0, 1, 1 + \zeta, 1, 0)$ and set $V_1 = \mathbf{F}v_1 \wedge v_2 + \mathbf{F}v_2 \wedge v_3 + \mathbf{F}v_1 \wedge v_3$, $V_2 = \mathbf{F}v_4 \wedge v_5$, and $V_3 = \mathbf{F}v_1 \wedge v_4 + \mathbf{F}v_1 \wedge v_5 + \mathbf{F}v_2 \wedge v_4 + \mathbf{F}v_2 \wedge v_5 + \mathbf{F}v_3 \wedge v_4 + \mathbf{F}v_3 \wedge v_5$. Then, we get the decomposition $\wedge^2 V = V_1 \oplus V_2 \oplus V_3$, and by Lemma 4.1, $\pi_\wp(\theta_1)$ acts on V_1, V_2, V_3 by the scalar multiples $\eta^{-4}, \eta^6, -\eta$, respectively, from which we see that $\wedge^2 W$ sits in one of V_i 's. Suppose $W = \mathbf{F}w \subset V_1$. Then, $\wedge^2 \pi(\theta_j)w$, $2 \leq j \leq 5$, should be in V_1 . Using the above base of V_1 and the assumption on \wp , just write down these and we get $w = 0$. Similarly, W can't be in V_2, V_3 . We conclude π_\wp is irreducible. \square

Now, we shall determine the image of ρ_p and there is a list of irreducible subgroups of $PSL_5(\mathbf{F})$ due to Martino and Wagoner [M-W]. Here, we assume further that p is prime to 2. By abuse of notation we write ρ_p for the associated projective representation and set $G = \rho_p(\Gamma)$, which is an irreducible subgroup of $PSL_5(\mathbf{F})$ by Lemma 3.2.

First, we have the following

Lemma 3.3. *The group G can not be realized over \mathbf{F}_{p^a} , $a < 2f$, where p^{2f} is the cardinality of \mathbf{F}*

Proof. Suppose that G is a subgroup of $PSL_5(\mathbf{F}_{p^a})$, $a < 2f$. Then, the characteristic polynomial $(X - \eta^{-2})(X + \eta^3)$ of $\rho_p(\theta_1)$ is invariant under the action of the Galois group $\text{Gal}(\mathbf{F}_{p^{2f}}/\mathbf{F}_{p^a}) = \langle \sigma \rangle$, where $\sigma =$ Frobenius automorphism, and so $\eta^\sigma = \eta^{p^a}$, by $(\eta^{-2})^\sigma = \eta^{-2}$. Since $(n, 10) = 1$, $p^a \equiv 1 \pmod{n}$. This contradicts to the minimality of $2f$ so that $p^{2f} \equiv 1 \pmod{n}$. \square

By Lemma 3.2, the following groups in the list of Martino-Wagoner can not be G : (1.3)-(a), (1.5), (1.7), (1.10)-(a), (1.12), (1.13), (1.14)-(a), (1.15), (1.16), where the numbers are those in [M-W].

Next, since the image of ρ_p is contained in $SU_5(\mathcal{O}_K/\mathfrak{p}_K; h_p) \simeq SU_5(\mathbf{F})$, G can not be $PSL_5(\mathbf{F})$, $PSO_5(\mathbf{F})$ and $P\Omega_5(\mathbf{F})$, by comparing the orders. So, the groups (1.4), (1.8), (1.9) and (1.10)-(b) in [M-W] are excluded.

The following useful lemma was suggested by Eiichi Bannai.

Lemma 3.4. *The subgroup of G generated by $\rho_p(\theta_1)$ and $\rho_p(\theta_3)$ is isomorphic to $\mathbf{Z}/2n\mathbf{Z} \times \mathbf{Z}/2n\mathbf{Z}$.*

Proof. By Lemma 3.1, the order of $\rho_p(\theta_i)$ is $2n$. We easily check $\langle \rho_p(\theta_1) \rangle \cap \langle \rho_p(\theta_3) \rangle = \text{id}$. \square

The group (1.2) in [M-W] is a subgroup of the group which is an extension of a cyclic subgroup by $\mathbf{Z}/5\mathbf{Z}$. So, by Lemma 3.4, G can not be this group. Next, (1.11) is $PSL_2(\mathbf{F})$ or $PGL_2(\mathbf{F})$. We have a list of subgroups of $PSL_2(\mathbf{F})$ due to Dickson, [H], p213, Satz 8.27. Looking at this, by Lemma 3.3, G can not be a subgroup of $PSL_2(\mathbf{F})$. Since $PGL_2(\mathbf{F})$ is an extension

of $PSL_2(\mathbf{F})$ by a cyclic subgroup of order 2, G can't be in $PGL_2(\mathbf{F})$. The similar argument can be applied to the groups (1.3)-(b),(c).

Finally, the group (1.1) can be excluded as follows (E. Bannai). The group (1.1) is an irreducible subgroup of A , where A is a global stabilizer in $PSL_5(\mathbf{F})$ of a simplex. Note that A is a monomial group and has a normal subgroup N so that $A/N \simeq S_5$ = the symmetric group on 5 letters. Assume that G is an irreducible subgroup of A . Then, $\bar{G} = G/(G \cap N)$ is a subgroup of S_5 and then \bar{G} can be one of S_5, A_5 , Frobenius group of order 20, dihedral group of order 10, or cyclic group of order 5. The images of $\rho_p(\theta_i)$ in \bar{G} satisfy the relation induced from that of the mapping class group, from which we can conclude \bar{G} is cyclic of order 5. This is a contradiction by the assumption $(n, 10) = 1$.

Summing up the above, we have

Theorem 3.5. *Assume that n is prime to 10, bigger than 2 and that a prime ideal \wp of \mathcal{O}_F does not divide $2(1 + \zeta)(\zeta + \zeta^{-1})(1 + \zeta + \zeta^{-1})$ and is inert in K/F . Then, the image of ρ_\wp coincides with $SU_5(\mathcal{O}_K/\wp_K; h_\wp)$.*

4. The case of a product of non-split primes

In this section, we extend Theorem 3.5 to the case where I is a product of non-split primes. For this, we apply a criterion of Weisfeiler on the approximation of a Zariski-dense subgroup in a semisimple group over a finite ring to our situation. In the following, we simply call (i) ~ (iv) for Weisfeiler's assumptions (i) ~ (iv) in (7.1) of [W].

Let I be a product of different prime ideals \wp_i of \mathcal{O}_F , $I = \prod_{i=1}^r \wp_i^{e_i}$, where each \wp_i is inert in K/F and prime to $6(1 + \zeta)(\zeta + \zeta^{-1})(1 + \zeta + \zeta^{-1})$. Set $A = \mathcal{O}_F/I$ and $B = \mathcal{O}_K/I_K$, $I_K = I\mathcal{O}_K$. Write $\mathbf{F}_{q_i} = \mathcal{O}_F/\wp_i$, $q_i = N\wp_i$, for simplicity. The radical of A is $R = \prod_{i=1}^r \wp_i$.

Let G_h and G be the special unitary group schemes over A with respect to the hermitian forms $h_I = h_\zeta \bmod I_K$ and $1_5 \in M_5(B)$ on the free B -module $M = B^{\oplus 5}$, respectively.

Our task is to show $G_h(A) = \rho_I(\Gamma)$. Fixing an isometry $\phi : (M; h_I) \simeq (M; 1_5)$ of hermitian modules, it is reduced to show $G(A) = \Gamma'$, where $\Gamma' = \phi \rho_I(\Gamma) \phi^{-1}$.

Let T_1 be the norm 1 torus attached to the quadratic extension B/A ;

$T_1 := \text{Ker}(R_{B/A}(\mathbf{G}_{\mathbf{m},B}) \xrightarrow{N} \mathbf{G}_{\mathbf{m},A})$, where $\mathbf{G}_{\mathbf{m}}$ is the split multiplicative group scheme of dimension 1 and $R_{B/A}$ is the Weil restriction of the scalar, and N is the norm map attached to B/A .

A maximal A -torus of G is given by $T := \{t = \text{diag}(t_1, t_2, t_3, t_4, t_5) \mid t_i \in T_1, \prod_{i=1}^5 t_i = 1\}$. Fix an isomorphism $T_1 \simeq \mathbf{G}_{\mathbf{m}}$ over B and define the character χ_i of T by $\chi_i(t) := t_i$, $1 \leq i \leq 4$. Then, the character module $X^*(T)$ of T is generated by χ_i , $1 \leq i \leq 4$. Suppose that $\chi|_{T(\mathbf{F}_{q_i})} = \chi'|_{T(\mathbf{F}_{q_i})}$ for $\chi, \chi' \in X^*(T)$. Then, writing χ and χ' as products of powers of χ_i 's, we easily see that $\chi = \chi'$. So, the assumption (i) is just $q_i \geq 10$, $1 \leq i \leq r$. The assumption (ii) is satisfied for our G and (iii) is a consequence of Theorem 3.5.

Finally, let $\text{Ad} : G(A) \rightarrow GL(L(A))$ be the adjoint representation, where L is the Lie algebra of G and given by $L(A) = \{X \in M_5(B) \mid \text{tr}(X) = 0, {}^tX^\sigma + X = 0\}$. Write $B = A + A\beta$, $\beta^2 \in A$, and take $\beta(e_{11} - e_{55}), \dots, \beta(e_{44} - e_{55}), e_{ij} - e_{ji}, \dots, \beta(e_{ij} + e_{ji})$, ($i < j$) as a basis of $L(A)$. Using this basis, a straightforward calculation shows that $\text{tr}(\text{Ad}(g)) = N_{B/A}(\text{tr}(g)) - 1$ for $g \in G(A)$, where $N_{B/A}$ is the Norm map attached to B/A and $N_{B/A}(\text{tr}(\rho_I(\theta_1))) = 13 - 6(\zeta + \zeta^{-1})$. From this, we get $\mathbf{Z}[\text{tr} \text{Ad}(\Gamma') \bmod R^2] = A/R^2$ which certifies the assumption (iv).

Summing up the above, we have

Main Theorem 4.1. *Let I be a product of prime ideals \wp_i of \mathcal{O}_F . Assume that each \wp_i is inert in K/F and prime to $6(1+\zeta)(\zeta+\zeta^{-1})(1+\zeta+\zeta^{-1})$ and $N_{\wp_i} \geq 10$. Then, the image of ρ_I coincides with $SU_5(\mathcal{O}_K/I_K, h_I)$.*

5. Comparison with the Torelli group and coverings of the moduli space of compact Riemann surfaces of genus 2

Let $Sp_2(\mathbf{Z})$ be the Siegel modular group of degree 4, namely, the group consisting of all $S \in GL_n(\mathbf{Z})$ such satisfying

$$SJ {}^tS = J, \quad J = \begin{pmatrix} 0_2 & 1_2 \\ -1_2 & 0_2 \end{pmatrix}.$$

Let $\theta : \Gamma \rightarrow Sp_2(\mathbf{Z})$ be a canonical homomorphism induced by the abelianization map of Γ and the Nielsen isomorphism. We call the kernel of θ the

Torelli group of genus 2 and write $\Gamma(N)$ for $\theta^{-1}(Sp_2(\mathbf{Z}; N))$, where $Sp_2(\mathbf{Z}; N)$ is the principal congruence subgroup of $Sp_2(\mathbf{Z})$ modulo a natural number N . The following result of Birman allows us to compare our groups $\Gamma_{n,I}$ with the Torelli group and $\Gamma(N)$.

Lemma 5.1. ([Bi1], Theorem 2) *The Torelli group of genus 2 is generated by the normal closure of $(\theta_1\theta_2\theta_1)^4$.*

Proposition 5.2. *Under the same assumption in Theorem 4.1, the group $\Gamma_{n,I}$ does not contain the Torelli group, hence any $\Gamma(N)$.*

Proof. It is straightforward to check that $\rho_{n,I}((\theta_1\theta_2\theta_1)^4) \neq 1$. \square

The geometrical interpretation of the above result is as follows.

Let \mathcal{T} be the Teichmüller space of genus 2 and $\mathcal{M} = \mathcal{T}/\Gamma$ be the moduli space of compact Riemann surfaces of genus 2. Let \mathcal{S} be the Siegel upper half space of degree 4 and $\mathcal{A} = \mathcal{S}/Sp_2(\mathbf{Z})$ be the moduli space of principally polarized abelian varieties. The period map $\mathcal{T} \rightarrow \mathcal{S}$ is compatible with the actions of Γ , $Sp_2(\mathbf{Z})$ and θ , and thus we obtain the Torelli map $\mathcal{M} \rightarrow \mathcal{A}$.

The Galois covering $\mathcal{A}_N = \mathcal{S}/Sp_2(\mathbf{Z}; N)$ over \mathcal{A} with the Galois group $Sp_2(\mathbf{Z}/N\mathbf{Z})$ is the moduli space of principally polarized abelian varieties with level N -structure. Then, Corollary 5.2 tells us that the spaces $\mathcal{T}/\Gamma_{n,I}$ give a family of Galois coverings over \mathcal{M} with the Galois groups $SU_5(\mathcal{O}_K/I_K)$, which can not be obtained by the pull-back of any \mathcal{A}_N via the Torelli map.

Acknowledgement. I would like to thank Takayuki Oda for explaining his conjecture and problems related to the moduli space of curves and useful discussions. My thanks also go to Eiichi Bannai for supplying some ideas and proofs in Section 3. A part of this work was done while I stayed at RIMS, Kyoto University, in the fall of 1995. It is my pleasure to thank Professor Yasutaka Ihara for giving me the opportunity to join his friendly Number Theory Seminar.

References

[Be] G. Berger, Fake congruence modular curves and subgroups of the modular group, (1995), preprint.

- [Bi1] J. Birman, On Siegel modular group, Math. Ann., **191**, (1971), 59-68.
- [Bi2] J. Birman, Braids, links and mapping class groups, Ann. Math. Studies, **82**, (1974).
- [J] V.F.R. Jones, Hecke algebra representations of braid groups and link polynomials, Ann. of Math., **126** (1987), 335-388.
- [H] Huppert, Endliche Gruppen I, Glundl. der math. Wiss. **134**, Springer (1967).
- [M-W] L.D. Martino and A. Wagoner, The irreducible subgroups of $PSL(V_5, q)$, where q is odd, Resultate d. Math. **2**. (1978).
- [O-T] T. Oda and T. Terasoma, Surjectivity of reduction of the Burau representations of Artin braid groups, in preparation (1996)
- [W] B. Weisfeiler, Strong approximation for Zariski-dense subgroups of semisimple algebraic groups, Ann. of Math., **120**, (1984), 271-315